

# Interlude

Yadav Munde<sup>1</sup>

<sup>1</sup> Ruby Hall Clinic, Pune

J Clin Interv Radiol 2016;1:31.


**Address for correspondence** Yadav Munde, MBBS, DNB, FVIR, Consultant Vascular and Interventional Radiologist, Flat #1003, F2-Building, Ganga Skies, Vallabh Nagar, Pimpri, Pune 411018, Maharashtra, India (e-mail: yadavmunde@gmail.com).

## Image description


A 39-year-old male patient with a history of road traffic accident presented with swelling (►**Fig. 1**) in the right calf. The digital subtraction angiography (►**Figs. 2, 3**) reveals a “heart symbol” shaped pseudoaneurysm arising from the proximal right anterior tibial artery with a wide and short neck. The pseudoaneurysm was treated by placing a stent graft (3.5 × 27 mm; Prograft, Vascular Concepts) (►**Fig. 4**) across the neck of pseudoaneurysm.




**Fig. 1** Clinical photograph reveals upper right leg swelling.



**Fig. 2** Color image from digital subtraction angiography reveals a heart shaped pseudoaneurysm from right anterior tibial artery.



**Fig. 3** Digital subtraction angiography of right leg demonstrating a heart shaped pseudoaneurysm from right anterior tibial artery.



**Fig. 4** Digital subtraction angiography following the exclusion of the pseudoaneurysm with a stent graft demonstrates patent right anterior tibial artery with no opacification of the pseudoaneurysm.